



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 471/04, A61K 31/435, 31/505, 31/425, C07D 487/04, 513/04 // (C07D 471/04, 221:00, 209:00) (C07D 471/04, 235:00, 221:00) (C07D 487/04, 239:00, 235:00) (C07D 513/04, 277:00, 235:00)</p>	A1	<p>(11) International Publication Number: WO 98/18797</p> <p>(43) International Publication Date: 7 May 1998 (07.05.98)</p>
<p>(21) International Application Number: PCT/HU97/00069</p> <p>(22) International Filing Date: 22 October 1997 (22.10.97)</p> <p>(30) Priority Data: P 96 03001 30 October 1996 (30.10.96) HU</p> <p>(71) Applicant (for all designated States except US): RICHTER GEDEON VEGYÉSZETI GYÁR RT. [HU/HU]; Gyömrői út 19.-21, H-1103 Budapest X (HU).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): LASZLOVSZKY, István [HU/HU]; Bartók Béla út 16, H-1111 Budapest (HU). DOMÁNY, György [HU/HU]; Bimbó u. 114/a, H-1022 Budapest (HU). ACS, Tibor [HU/HU]; Szilárd u. 92, H-1174 Budapest (HU). FERENCZY, György [HU/HU]; Orbánhegyi út 23, H-1126 Budapest (HU). SZÁNTAY, Csaba, Jr [HU/HU]; Zsombolyai u. 8/2, H-1113 Budapest (HU). THUROCCZY, KÁLMÁN, Eszter [HU/HU]; Törökugrató út 5. I, H-1118 Budapest (HU). LAPIS, Erzsébet [HU/HU]; Abaliget u. 86, H-1172 Budapest (HU). TRISCHLER, Ferenc [HU/HU]; Úttörő u. 16, H-1171 Bu-</p>		<p>dapest (HU). HEGEDÚS, Béla [HU/HU]; Bartók Béla út 82, H-1113 Budapest (HU). AUTH, Ferenc [HU/HU]; Péteri u. 218, H-1188 Budapest (HU). CSEJTEI, Mónika [HU/HU]; Kós Károly tér 15, H-1192 Budapest (HU). KÁRPÁTI, Egon [HU/HU]; Mihályfi Ernő u. 7/b, H-1022 Budapest (HU). KISS, Béla [HU/HU]; Gergely u. 48, H-1103 Budapest (HU). LASZY, Judit [HU/HU]; Beregszász u. 40, H-1112 Budapest (HU). PELLIONISZNÉ PARÓCZAI, Margit [HU/HU]; Trencsényi út 45, H-1125 Budapest (HU). SARKADI, Ádám [HU/HU]; Thököly út 85, H-1146 Budapest (HU). SZABÓ, Sándor [HU/HU]; Liszt Ferenc tér 9, H-1061 Budapest (HU).</p> <p>(74) Common Representative: RICHTER GEDEON VEGYÉSZETI GYÁR RT.; Gyömrői út 19.-21., H-1103 Budapest X (HU).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>
<p>(54) Title: 2-METHOXYPHENYLPIPERAZINE DERIVATIVES</p> <div style="text-align: center;"> <p>(I)</p> </div> <p>(57) Abstract</p> <p>The invention relates to novel fused heterocyclic compounds with ring junction nitrogen atom of formula (I), wherein Q stands for 2-indolizinyl, 2-imidazo[1,2-a]pyridinyl, 2-imidazo[1,2-a]pyrimidinyl, 6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl or 6-imidazo[2,1-b]thiazolyl group; and n is an integer from 2 to 4, as well as their therapeutically useful salts thereof. The invention further relates to pharmaceutical compositions containing these compounds as well as a process for the preparation of the above compounds and compositions. The novel compounds of formula (I) exhibit mainly antipsychotic effects so the invention relates also to a method of treatment of schizophrenia, organic mental disorders, affective disorders, anxiety and personality disorders.</p>		


FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

2-METHOXYPHENYLPIPERAZINE DERIVATIVES

The invention relates to novel fused heterocyclic compounds with ring junction nitrogen atom of a general formula (I)




(I)

wherein

Q stands for 2-indolizinyl, 2-imidazo[1,2-a]pyridinyl, 2-imidazo[1,2-a]-pyrimidinyl, 6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl or 6-imidazo[2,1-b]-thiazolyl group; and


n is an integer from 2 to 4,

and to their therapeutically useful salts and pharmaceutical compositions containing these compounds. Furthermore, the invention relates also to a process for the preparation of the compounds of formula (I) and their therapeutically useful salts in such a way that a haloalkyl ether derivative of formula (II)



(II)

wherein Q and n are as defined above, and X means halogen is reacted with (2-methoxyphenyl)piperazine of formula (III).



(III)

The compounds of formula (I) according to the invention are novel and possess significant biological activity, first of all antipsychotic effects.

5 The invention relates also to a method of treatment, which comprises administering a therapeutically active amount of a compound of formula (I) or a therapeutically acceptable salt thereof to a patient to be treated.

Among the starting substances, some of chloroalkyl ether derivatives of formula (II) are known from the literature, such as 3-[4-(2-imidazo[1,2-
10 a]pyridinyl)-phenoxy]propyl chlorid, 3-[4-(2-imidazo[1,2-a]pyrimidinyl)phenoxy]propyl chloride, or 3-[4-(6-imidazo[2,1-b]thiazolyl)phenoxy]propyl chloride [J. Med. Chem. 31, 2221 (1988)]. Other starting substances of formula (II) can be obtained according to the preparation process described in the above-cited literature.

15 (2-Methoxyphenyl)piperazine of formula (III) is a known, commercially available substance.

Compounds structurally similar to the substances of formula (I) are known from the literature. Such (substituted amino)propoxyphenylimidazo[1,2-
20 a]pyridines, substituted amino)propoxyphenylimidazo[1,2-a]pyrimidines and (substituted amino)propoxyphenylimidazo[2,1-b]thiazoles are described in the above-cited publication [J. Med. Chem. 31, 2221 (1988)], however, unlike the compounds according to the invention, these compounds possess calcium channel-blocking and local anaesthetic effects.

25 In contrast to the structurally closely related compounds known from the literature, the novel compounds of formula (I) according to the invention are orally effective and are endowed of a considerable neuroleptic activity. Based

on their biological activity, these compounds can be used as atypical antipsychotics, antidepressants, anxiolytics, neuroprotective and/or cognitive function improving agents, antiemetics or antiaddictives.

Since the 70's, antipsychotics have successfully been employed for the treatment of schizophrenia. Up to the present haloperidol has been used in the clinical practice most widely. Phenothiazines and haloperidol played a pioneering role in this therapeutic field and considerably contributed to the development of the dopamine theory of schizophrenia. Later researches confirmed also the role of serotonin and a number of other neurotransmitter systems, such as histaminerg, α -adrenerg, CCK-erg, etc. in this disorder. However, haloperidol and other typical antipsychotics improve only the positive symptoms of the disease, e. g. hallucinations, delusions, agitation and thought disturbances, whereas negative symptoms, such as emotional obtusion, autism, social isolation, neglect of personal hygiene remain unimproved. In addition, nearly 30% of the patients do not respond to the treatment and a number of undesired adverse effects cannot be excluded, either. From these, the most severe adverse effects are the appearance of extrapyramidal symptoms (EPS), because of the strong but not regionselective antagonism of D-2 dopamine receptor, deterioration of cognitive functions caused by anticholinergic effect, orthostatic hypotonia (α -adrenerg antagonism), and hyperprolactinaemia.

Nowadays, intensive research is being devoted to atypical antipsychotics, which are capable of improving both positive and negative symptoms, do not induce extrapyramidal symptoms or only in doses higher than therapeutical, only a few non-responsive patients should be taken into account at all if any and other side effects are also negligible. Clozapine, if did not cause agranulocytosis, could be such an ideal antipsychotic.

Nowadays research is being focused on discovering clozapine-like atypical antipsychotics. Clozapine-like action means that the molecule possesses a strong antipsychotic effect without appearance of the above side effects. Such compounds have a direct or indirect selective action on the

dopaminergic pathways of limbic brain and this effect is associated with a complex receptor profile. Within the complex receptor profile, the D-2 receptor antagonistic effect characteristic of haloperidol is not dominant.

The novel compounds of formula (I) according to the invention exert an antipsychotic effect similar to that of the atypical antipsychotic clozapin. Both antipsychotic and oral activities of the compounds were confirmed by apomorphine-induced climbing and sniffing tests *in vivo*. The mechanism of action of the compounds was characterized by receptor binding tests *in vitro*.

Inhibition of apomorphine (APO) induced climbing and sniffing

CD-1 (Charles River) male mice weighing 22 to 24 g were pretreated with 1% Tween 80 solution of the compound to be tested. After 55 minutes, the animals were placed two by two into examination cages [P. Potrais et al.: Psychopharmacol 50, 1-6 (1976)]. Six animals were used in each group. At 60 minutes after pretreatment, the animals were treated subcutaneously with 1 mg/kg of apomorphine (APO). From the 10th to 25th minute following APO treatment the animals were scored in each minute as follows: 0: all the four legs of the animal are on the ground; 1: the animal climbs on the grid by its both forelegs; 2: the animal climbs on the grid by all its four legs. The inhibitory effect was related to the APO control group, which could achieve 32 scores as a maximum.

The inhibition of APO-induced sniffing stereotypy was measured simultaneously with the inhibition of climbing from the 10th to 25th minute following the APO administration according to S. Gerhardt [S. Gerhardt et al.: Life Sci. 37, 2355-2363 (1985)].

Receptor binding assays

D-2

Binding to the D-2 dopamine receptor was studied according to P. Seeman [P. Seeman et al.: J. Neurochem. 43, 221-235 (1984)] by using 0.5 nM

³ H-spiperone as ligand; the non-specific binding was determined in the presence of 10 μ M (\pm)-sulpiride.

5-HT_{1A}

Determination of the 5-HT_{1A} subtype of serotonin receptor was carried out by a modification [M.D. Hall et al.: J. Neurochem. 44, 1685-1696 (1985); H. Gozlan et al.: Nature 305, 140-142 (1983)] of the method described by Peroutka [S. J. Peroutka: J. Neurochem. 47, 529-540 (1986)]. 0.5 nM ³ H-8-OH-DPAT was used as ligand, whereas non-specific binding was determined by using 10 μ M serotonin.

Alpha-1

The α -1 receptor activity of the compounds was measured according to the method of Greengrass and Horung [P. Greengrass et al.: Eur. J. Pharmacol. 55, 323-326 (1979); R. Horung et al.: Naunyn-Schmiedeb. Arch Pharmacol. 308, 223-230 (1979)] by using ³ H-prazosin as ligand. The non-specific binding was measured with 10 μ M (\pm)-phentolamine.

The results of the in vivo and in vitro investigations are summarized in Table 1, where

Q: A means 2-indolizinyI,

Q: B means 2-imidazo[1,2-a]pyridinyl,

Q: C means 2-imidazo[1,2-a]pyrimidinyl,

Q: D means 6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl and

Q: E means 6-imidazo[2,1-b]thiazolyl group.

Table 1

Compound No.	Q	n	ED ₅₀ (mg/kg p.o.)		Sniff./ Climb.	Receptor IC ₅₀ (nM)		
			Inhibition of APO- induced climbing	Inhibition of APO- induced sniffing		D-2	5-HT _{1A}	α-1
4510407	A	2	» 30	-	-	258	802	542
4510423	A	3	4.1	> 10	>2.4	77	1000	359
4510613	A	4	14.8	> 30	>2.0	99	243	842
4510408	B	2	2.4	≈ 10	4.1	56	70	36
4510067	B	3	7.2	16.2	2.3	17	202	57
4510915	B	4	10.7	-	-	69	21	34
4510411	C	2	3.4	» 30	>8.8	133	10	63
4510424	C	3	8.9	» 30	3.4	34	66	30
4510645	C	4	20.5	» 30	>1.5	55	12	22
4510471	D	2	6.3	» 30	4.8	117	29	75
4510473	D	3	15.3	> 30	>2.0	29	122	48
4510991	D	4	≈ 30	» 30	>1.0	56	9	16
4510470	E	2	0.8	4.5	5.6	73	38	101
4510472	E	3	3.4	≈ 10	2.9	23	96	56
4510916	E	4	6.1	≈ 30	>4.9	55	8	28
haloperidol			0.19	0.4	2.1	3	3860	19
clozapine			3.3	12.3	3.7	159	647	35

It is evident from the ED₅₀ values of inhibition of the APO-induced climbing inhibition shown in Table 1 that most of the compounds inhibit the APO-induced climbing already in a low oral dose (0.8-10 mg/kg), a result supporting the in vivo antipsychotic activity of the compounds. The most active compound No. 4510470 is of ten-fold oral activity as compared to clozapine, but several other molecules, such as the compounds Nos. 4510408, 4510411 or 4510472, have ED₅₀ values lower than or equal to that of clozapine. Simultaneously, the apomorphine-induced stereotypy measured by the occurrence of sniffing in this case was prevented only in significantly higher doses usually above 30 mg/kg of the compounds according to the invention. The above results, including the sniffing/climbing ratio, confirm the compounds' limbic selectivity in vivo and this selectivity is better than that of clozapine. Based on the limbic selectivity, the action of the compounds is similar to that of atypical antipsychotics, thus, the occurrence of extrapyramidal symptoms (EPS) cannot be expected or only in higher doses.

The mechanism of action based on results of receptor binding assays shown in Table 1. provide further support to the atypical antipsychotic nature of the compounds of the invention. Thus, their D-2 dopamine receptor activity is less marked than the effect of the typical antipsychotic haloperidol (3 nM) and indicates similarity rather to the atypical antipsychotic clozapine. However, the compounds according to the invention are somewhat more effective on this receptor subtype. Their clozapine-like α -adrenerg receptor activity provides further evidence for their atypical features. The compounds in general as well as their representatives, such as molecules No. 4510916 and No. 4510991, exert especially strong activity on 5-HT_{1A} receptor subtypes. This effect is significantly different from the weak 5-HT_{1A} receptor activity of clozapine and similarly, it is different from the strength of activity of other atypical antipsychotics which have been commercially available up to the present. This difference is even more markedly expressed by the D-2/5-HT_{1A} ratio. For example, this ratio is 0.2 for clozapine, whereas it is 1.9 for the compound No.

4510470 and is even 6.9 for the compound No. 4510916. The 5-HT_{1A} receptor activity indicates also an anxiolytic effect component of the compounds.

In summary: the novel compounds of the invention are atypical antipsychotics, which are essentially more active than clozapine after oral administration. Similarly to clozapine, they possess a complex receptor profile, however, receptor subtypes and their strength of activity involved participating in the mechanism of action of the compounds are different from those of clozapine. Thus, the compounds of the invention are endowed of other character and new sort of complex profile of mechanism of action. Therefore, the compounds according to the invention could be very effective in the treatment of acute and chronic schizophrenia; paranoia and other psychotic disturbances; organic mental disorders, such as delirium, dementia, withdrawal syndromes, addictions, mental retardation, tic disorders; affective disorders, such as mania, bipolar disorders, cyclothymia, dysthymia; anxiety disorders, including panic disorders, phobia, obsessive-compulsive disorders, generalized anxiety syndrome and personality disorders, such as compulsive, paranoid, schizoid, antisocial and any other disorders related to psychomotor agitation. The expected therapeutical doses of the compounds are between 0.01 and 50 mg/kg of body weight, once or in repeated subdoses daily, administered orally, intraperitoneally or subcutaneously route.

The preparation of the novel compounds of formula (I) according to the invention are hereinafter described in detail.

The haloalkyl ether derivatives of formula (II), preferably 2-chloroalkyl ether derivative known from the literature or prepared by known methods, are reacted with the similarly known (2-methoxyphenyl)piperazine or a salt thereof, preferably with the commercially available (2-methoxyphenyl)piperazine dihydrochloride in an organic protic solvent, e.g. in an alcohol or in any organic dipolar-aprotic solvent, such as an aliphatic ketone; or acetonitrile, dimethylformamide and the like, optionally in the presence of a base and an alkaline metal iodide, such as sodium iodide. Suitable bases are inorganic bases, e.g. potassium carbonate, sodium carbonate; or organic bases, e.g.

triethylamine. The reaction is carried out at the boiling point of the solvent used, for a reaction time of 5 to 10 hours. After evaporation of the reaction mixture containing the compound of formula (I), treating the dry residue with water and extracting the obtained mixture with a water-immiscible solvent, the crude target compound of formula (I) is obtained, which can be purified by recrystallization if necessary.

If desired, the compounds of formula (I) can be converted to their acid addition salts in a manner known per se.

The salt formation is accomplished in a known manner in an inert organic solvent or solvent mixture in such a way that the compound of formula (I) is dissolved in the selected solvent and subsequently, the appropriate acid is added in portions to the above solution until the mixture becomes strongly acidic (about pH 1 value). Alternatively, the salt formation can be performed by adding the solution of the acid in the calculated amount in the selected solvents to the above solution. Thereafter, the precipitated acid addition salt is separated from the reaction mixture in a suitable manner, e.g. by filtration.

The active ingredient of formula (I) can be transformed to pharmaceutical compositions by mixing it with nontoxic, inert solid or liquid carriers commonly used in the therapy for parenteral or enteral administration. E.g. water, gelatin, lactose, starch, pectin, magnesium stearate, stearic acid, talc, vegetable oils such as olive oil or peanut oil and the like are useful carriers. The active ingredient may be formulated in the form of usual pharmaceutical compositions, particularly in solid form, e.g. as tablets, dragées, capsules, pills, suppository and the like. The amount of the solid carrier may be varied within wide limits, preferably between about 25 mg and 1 g. These compositions may optionally contain the commonly used pharmaceutical auxiliaries (additives), e.g. preservatives, stabilizing, wetting, emulsifying agents or the like. The preparation of these compositions can be accomplished by common methods, e.g. by sieving, mixing, granulating and then compressing the components in the case of solid compositions. The compositions may be exposed to further usual operations, e.g. sterilization.

The invention is illustrated in detail by the following non-limiting examples.

Example 1

1-(2-Methoxyphenyl)-4-{2-[4-(2-indolizinyI)phenoxy]ethyl}piperazine

[Formula (I), Q = 2-indolizinyI, n=2]

A mixture containing 2.72 g (10 mmol) of 2-[4-(2-indolizinyI)phenoxy]ethyl chloride, 2.74 g (12 mmol) of (2-methoxyphenyl)piperazine dihydrochloride, 2.54 g (24 mmol) of sodium carbonate, 0.3 g (2 mmol) of anhydrous sodium iodide and 40 ml of methyl isobutyl ketone is boiled under reflux for 10 hours. After evaporating the solvent under reduced pressure, the residue is thoroughly triturated with 20 ml of water, then extracted with 80 ml of chloroform. The organic phase is twice washed with 15 ml of water each, dried over anhydrous sodium sulfate and the drying agent is filtered off. The filtrate is stirred with 1 g of charcoal and 0.5 g of aluminium oxide for 20 minutes, then filtered and evaporated to its half volume under reduced pressure. After adding 40 ml of ethanol to the residue, it is evaporated to about 15 ml under reduced pressure. The precipitated material is filtered and dried. The obtained 2.9 g crude product is dissolved in 30 ml of chloroform, 30 ml of ethanol are added then, the solution obtained is evaporated to about 15 ml under reduced pressure. After filtering and drying the precipitated substance, the title compound is obtained in a yield of 2.61 g (61%), m.p.: 159-161 °C.

Example 2

1-(2-Methoxyphenyl)-4-{3-[4-(2-indolizinyI)phenoxy]propyl}piperazine

[Formula (I), Q = 2-indolizinyI, n = 3]

The procedure described in Example 1 is followed, except that 3-[4-(2-indolizinyI)phenoxy]propyl chloride is used as starting material instead of 2-[4-(2-indolizinyI)phenoxy]ethyl chloride. The title compound melts at 157-158 °C.

Example 3

1-(2-Methoxyphenyl)-4-{4-[4-(2-indolizinyI)phenoxy]butyl}piperazine

[Formula (I), Q = 2-indolizinyI, n = 4], compound No. 4510613

The procedure described in Example 1 is followed, except that 4-[4-(2-indolizinyloxy)phenoxy]butyl chloride is used as starting material instead of 2-[4-(2-indolizinyloxy)phenoxy]ethyl chloride. The title compound melts at 153-155 °C.

Example 4

5 **1-(2-Methoxyphenyl)-4-{2-[4-(2-imidazo[1,2-a]pyridinyl)phenoxy]ethyl}piperazine**

[Formula (I), Q= 2-imidazo[1,2-a]pyridinyl, n = 2], compound No. 4510408

The procedure described in Example 1 is followed, except that 2-[4-(2-imidazo[1,2-a]pyridinyl)phenoxy]ethyl chloride is used as starting material instead of 2-[4-(2-indolizinyloxy)phenoxy]ethyl chloride. The title compound melts at 151-153 °C.

Example 5

15 **1-(2-Methoxyphenyl)-4-{3-[4-(2-imidazo[1,2-a]pyridinyl)phenoxy]propyl}piperazine**

[Formula (I), Q= 2-imidazo[1,2-a]pyridinyl, n = 3], compound No. 4510067

The procedure described in Example 1 is followed, except that 3-[4-(2-imidazo[1,2-a]pyridinyl)phenoxy]propyl chloride is used as starting material instead of 2-[4-(2-indolizinyloxy)phenoxy]ethyl chloride. The title compound melts at 149-150 °C.

Example 6

20 **1-(2-Methoxyphenyl)-4-{4-[4-(2-imidazo[1,2-a]pyridinyl)phenoxy]butyl}piperazine**

[Formula (I), Q= 2-imidazo[1,2-a]pyridinyl, n = 4], compound No. 4510915

The procedure described in Example 1 is followed, except that 4-[4-(2-imidazo[1,2-a]pyridinyl)phenoxy]butyl chloride is used as starting material instead of 2-[4-(2-indolizinyloxy)phenoxy]ethyl chloride. The title compound melts at 144-145 °C.

Example 7

30 **1-(2-Methoxyphenyl)-4-{2-[4-(2-imidazo[1,2-a]pyrimidinyl)phenoxy]ethyl}piperazine**

[Formula (I), Q=2-imidazo[1,2-a]pyrimidinyl, n=2], compound No.4510911

The procedure described in Example 1 is followed, except that 2-[4-(2-imidazo[1,2-a]pyrimidinyl)phenoxy]ethyl chloride is used as starting material instead of 2-[4-(2-indoliziny]phenoxy]ethyl chloride. The title compound melts at 150-152 °C.

Example 8

1-(2-Methoxyphenyl)-4-{ 3-[4-(2-imidazo[1,2-a]pyrimidinyl)phenoxy]propyl}piperazine

[Formula(I), Q=2-imidazo[1,2-a]pyrimidinyl, n=3], compound No. 4510924

The procedure described in Example 1 is followed, except that 3-[4-(2-imidazo[1,2-a]pyrimidinyl)phenoxy]propyl chloride is used as starting material instead of 2-[4-(2-indoliziny]phenoxy]ethyl chloride. The title compound melts at 156-158 °C.

Example 9

1-(2-Methoxyphenyl)-4-{ 4-[4-(2-imidazo[1,2-a]pyrimidinyl)phenoxy]butyl}piperazine

[Formula(I), Q=2-imidazo[1,2-a]pyrimidinyl, n=4], compound No. 4510645

The procedure described in Example 1 is followed, except that 4-[4-(2-imidazo[1,2-a]pyrimidinyl)phenoxy]butyl chloride is used as starting material instead of 2-[4-(2-indoliziny]phenoxy]ethyl chloride. The title compound melts at 151-153 °C.

Example 10

1-(2-Methoxyphenyl)-4-{ 2-[4-[6-(2,3-dihydroimidazo[2,1-b]thiazol)yl]phenoxy]ethyl}piperazine

[Formula (I), Q = 6-(2,3-dihydroimidazo[2,1-b]thiazol)yl, n = 2],

The procedure described in Example 1 is followed, except that 2-{4-[6-(2,3-dihydroimidazo[2,1-b]thiazol)yl]phenoxy]ethyl} chlorid is used as starting material instead of 2-[4-(2-indoliziny]phenoxy]ethyl chloride. The title compound melts at 145-147 °C.

Example 11

1-(2-Methoxyphenyl)-4-{ 3-[4-[6-(2,3-dihydroimidazo[2,1-b]thiazol)yl]phenoxy]propyl}piperazine

[Formula (I), Q = 6-(2,3-dihydroimidazo[2,1-b]thiazol)yl, n = 3], compound No. 4510473

The procedure described in Example 1 is followed, except that 3-{4-[6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl]phenoxy}propyl chlorid is used as starting material instead of 2-[4-(2-indoliziny]phenoxy]ethyl chloride. The title compound melts at 142-144 °C.

Example 12

1-(2-Methoxyphenyl)-4-{ 4-[4-[6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl]phenoxy]butyl}piperazine

[Formula (I), Q = 6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl, n = 4], compound No. 4510991

The procedure described in Example 1 is followed, except that 4-{4-[6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl]phenoxy}butyl chlorid is used as starting material instead of 2-[4-(2-indoliziny]phenoxy]ethyl chloride. The title compound melts at 139-140 °C.

Example 13

1-(2-Methoxyphenyl)-4-{ 2-[4-[6-(imidazo[2,1-b]thiazolyl)phenoxy]ethyl}piperazine

[Formula (I), Q = 6-imidazo[2,1-b]thiazolyl, n = 2],

The procedure described in Example 1 is followed, except that 2-{4-[6-imidazo[2,1-b]thiazolyl]phenoxy}ethyl chlorid is used as starting material instead of 2-[4-(2-indoliziny]phenoxy]ethyl chloride. The title compound melts at 158-161 °C.

Example 14

1-(2-Methoxyphenyl)-4-{ 3-[4-[6-imidazo[2,1-b]thiazolyl)phenoxy]propyl}piperazine

[Formula (I), Q = 6-imidazo[2,1-b]thiazolyl, n = 3], compound No. 4510472

The procedure described in Example 1 is followed, except that 3-[4-(6-imidazo[2,1-b]thiazolyl)phenoxy]propyl chloride is used as starting material instead of 2-[4-(2-indoliziny)phenoxy]ethyl chloride. The title compound melts at 130-132 °C.

Example 15

1-(2-Methoxyphenyl)-4-{4-[4-[6-imidazo[2,1-b]thiazolyl)phenoxy]butyl}piperazine

[Formula (I), Q = 6-imidazo[2,1-b]thiazolyl, n = 4]

The procedure described in Example 1 is followed, except that 4-[4-[6-imidazo[2,1-b]thiazolyl)phenoxy]butyl chloride is used as starting material instead of 2-[4-(2-indoliziny)phenoxy]ethyl chloride. The title compound melts at 128-130 °C.

Example 16

1-(2-Methoxyphenyl)-4-{2-[4-(2-indoliziny)phenoxy]ethyl}piperazine dihydrochloride

[Formula (I), Q = 2-indoliziny, n = 2], compound No. 4510407

After dissolving 21.5 g (0.05 mol) of 1-(2-methoxyphenyl)-4-{2-[4-(2-indoliziny)phenoxy]ethyl}piperazine (prepared as described in Example 1) in 50 ml of chloroform, 50 ml of ethanol are added to the solution, then the reaction mixture is acidified to pH 1 by adding 20 to 30% ethanolic hydrochloric acid. The precipitate is filtered, washed with a little amount of ethanol and dried to yield 24.5 g (98%) of title compound, m.p. 220-222 °C.

Example 17

1-(2-Methoxyphenyl)-4-{3-[4-[2-indoliziny)phenoxy]propyl}piperazine dihydrochloride

[Formula (I), Q = 2-indoliziny, n = 3], compound No. 4510423

The procedure described in Example 16 is followed, except that 1-(2-methoxyphenyl)-4-{3-[4-(2-indoliziny)phenoxy]propyl}piperazine is used as starting substance instead of 1-(2-methoxyphenyl)-4-{2-[4-(2-indoliziny)phenoxy]ethyl}piperazine. The title compound melts at 239-241 °C.

Exempl 18

1-(2-Methoxyphenyl)-4-{2-[4-[6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl]phenoxy]ethyl}piperazine trihydrochloride

[Formula (I), Q = 6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl, n=2], compound No. 4510471

The procedure described in Example 16 is followed, except that 1-(2-methoxyphenyl)-4-{2-[4-(6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl]phenoxy}ethyl piperazine is used as starting substance instead of 1-(2-methoxyphenyl)-4-{2-[4-(2-indoliziny)]phenoxy}ethyl}piperazine. The title compound melts at 214-216°C.

Example 19

1-(2-Methoxyphenyl)-4-{2-[4-[6-imidazo[2,1-b]thiazolyl]phenoxy]ethyl} piperazine trihydrochloride

[Formula (I), Q=6-imidazo[2,1-b]thiazolyl, n = 2], compound No. 4510470

The procedure described in Example 16 is followed, except that 1-(2-methoxyphenyl)-4-{2-[4-(6-imidazol[2,1-b]thiazolyl]phenoxy]ethyl}piperazine is used as starting material instead of 1-(2-methoxyphenyl)-4-{2-[4-(2-indoliziny)]phenoxy] ethyl}piperazine. The title compound melts at 254-256°C.

Example 20


1-(2-Methoxyphenyl)-4-{4-[4-[6-imidazo[2,1-b]thiazolyl]phenoxy]butyl} piperazine trihydrochloride

[Formula (I), Q=6-imidazo[2,1-b]thiazolyl, n = 4], compound No. 4510916

The procedure described in Example 16 is followed, except that 1-(2-methoxyphenyl)-4-{4-[4-(6-imidazol[2,1-b]thiazolyl]phenoxy]butyl}piperazine is used as starting material instead of 1-(2-methoxyphenyl)-4-{2-[4-(2-indoliziny)]phenoxy] ethyl}piperazine. The title compound melts at 227-229°C.

Claims

1. Novel fused heterocyclic compounds with ring junction nitrogen atom of formula (I)



(I)


wherein

Q stands for 2-indolizinyl, 2-imidazo[1,2-a]pyridinyl, 2-imidazo[1,2-a]pyrimidinyl, 6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl or 6-imidazo[2,1-b]thiazolyl group and

n is an integer of 2 to 4,
as well as their therapeutically acceptable salts.

2. The compounds of formula (I) according to claim 1, wherein Q stands for 2-imidazo[1,2-a]pyridinyl or 6-imidazo[2,1-b]thiazolyl group; and n is as defined in claim 1, as well as their therapeutically acceptable salts.

3. Pharmaceutical compositions which comprises as active ingredient a compound of formula (I),



(I)


wherein

Q stands for 2-indolizinyl, 2-imidazo[1,2-a]pyridinyl, 2-imidazo[1,2-a]pyrimidinyl, 6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl or 6-imidazo[2,1-b]thiazolyl group and

n is an integer of 2 to 4

or a therapeutically acceptable salt thereof in admixture with a carrier and/or other additives commonly used in the pharmaceutical industry.

4. Process for the preparation of the novel fused heterocyclic compounds with ring junction nitrogen atom of formula (I),




(I)

wherein

Q stands for 2-indolizinyl, 2-imidazo[1,2-a]pyridinyl, 2-imidazo[1,2-a]pyrimidinyl, 6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl or 6-imidazo[2,1-b]thiazolyl group; and


n is an integer of 2 to 4,

as well as therapeutically acceptable salts thereof, which comprises reacting a haloalkyl ether derivative of formula (II)



(II)

wherein Q and n are as defined above, and X means halogen with (2-methoxyphenyl)piperazine of formula (III)



(III)


5

or a salt thereof in an organic solvent; and if desired converting a compound of formula (I) to therapeutically acceptable salt thereof.

5. A process according to claim 4, which comprises reacting the compound of formula (II) with the compound of formula (III) in the presence of base and alkaline metal iodide.

10

6. Process for the preparation of a pharmaceutical composition, which comprises mixing as active ingredient a novel fused heterocyclic compound with ring junction nitrogen atom of formula (I),



(I)

15


wherein

Q stands for 2-indolizinyl, 2-imidazo[1,2-a]pyridinyl, 2-imidazo[1,2-a]pyrimidinyl, 6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl or 6-imidazo[2,1-b]thiazolyl group, and

n is an integer of 2 to 4,

or a therapeutically acceptable salt thereof with a carrier and/or other additives commonly used in the pharmaceutical industry, and converting the mixture to a pharmaceutical composition.

7. Method of treatment of schizophrenia, organic mental disorders, affective disorders, anxiety and personality disorders, w h i c h comprises administering to a patient to be treated a therapeutically effective amount of a novel fused heterocyclic compound with ring junction nitrogen atom of formula (I),



(I)


wherein

Q stands for 2-indolizinyl, 2-imidazo[1,2-a]pyridinyl, 2-imidazo[1,2-a]pyrimidinyl, 6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl or 6-imidazo[2,1-b]thiazolyl group and

n is an integer of 2 to 4,

or a therapeutically acceptable salt thereof alone or in the form of a pharmaceutical composition.

8. Use of a compound of formula (I)



(I)


wherein

Q stands for 2-indolizinyl, 2-imidazo[1,2-a]pyridinyl, 2-imidazo[1,2-a]-
pyrimidinyl, 6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl or 6-imidazo[2,1-b]thiazolyl
group and

n is an integer of 2 to 4,

or a therapeutically acceptable salt thereof for the preparation of
pharmaceutical compositions for the treatment of schizophrenia, organic mental
disorders, affective disorders, anxiety and personality disorders.

9. Use of a compound of formula (I)



(I)

wherein

Q stands for 2-indolizinyl, 2-imidazo[1,2-a]pyridinyl, 2-imidazo[1,2-a]-
pyrimidinyl, 6-(2,3-dihydroimidazo[2,1-b]thiazol)-yl or 6-imidazo[2,1-b]thiazolyl
group and

n is an integer of 2 to 4,

or a therapeutically acceptable salt thereof for the treatment of schizophrenia, organic mental disorders, affective disorders, anxiety and personality disorders.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/HU 97/00069

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D471/04 A61K31/435 A61K31/505 A61K31/425 C07D487/04
C07D513/04 // (C07D471/04, 221:00, 209:00), (C07D471/04, 235:00,
221:00), (C07D487/04, 239:00, 235:00), (C07D513/04, 277:00, 235:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 30659 A (WARNER-LAMBERT) 16 November 1995 see claims 1,4,5 -----	1,6

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

16 February 1998

Date of mailing of the international search report

05.03.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Alfaro Faus, I

INTERNATIONAL SEARCH REPORT

International application No.

PCT/HU 97/ 00069

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 7, 9
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 7 and 9
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/HU 97/00069

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9530659 A	16-11-95	US 5486517 A	23-01-96
		AU 2197895 A	29-11-95
		ZA 9503752 A	11-01-96
